


Heavy Neutrinos in Ice, Rock, Water, and Plastic

Nicholas Kamp | nkamp@g.harvard.edu

Universiteit Gent | EPPGA Seminar

4 February 2025



Conwaybreen Glacier, Svalbard



The *Big* Picture

The *Big* Picture

Progress in neutrino physics is driven by **experimental anomalies**

The *Big* Picture

Progress in neutrino physics is driven by **experimental anomalies**

Novel detectors and unique signatures are essential to resolve anomalies

The *Big* Picture

Progress in neutrino physics is driven by **experimental anomalies**

Novel detectors and unique signatures are essential to resolve anomalies

This talk: a new perspective on an old anomaly

A Brief History of Neutrino Anomalies

Experimental Anomaly


Theoretical Resolution

Experimental Validation

A Brief History of Neutrino Anomalies

Experimental Anomaly

Continuous electron energy spectrum in beta decays



Neary 1940


Theoretical Resolution

Experimental Validation

A Brief History of Neutrino Anomalies

Experimental Anomaly


Continuous electron energy spectrum in beta decays



Neary 1940

Theoretical Resolution

A new neutral particle




Pauli 1930

Experimental Validation

A Brief History of Neutrino Anomalies

Experimental Anomaly

Continuous electron energy spectrum in beta decays



Theoretical Resolution

A new neutral particle

Physikalisches Institut
der Eidg. Technischen Hochschule
Zürich

Zürich, 4. Dez. 1930
Gloriastrasse

Liebe Radioaktive Damen und Herren,
Wie der Ueberbringer dieser Zeilen, den ich huldvollst
ansuhören bitte, Ihnen des näheren auseinandersetzen wird, bin ich
angesichts der "falschen" Statistik der N - und Li-6 Kerne, sowie
des kontinuierlichen beta-Spektrums auf einen verswiefelten Ausweg
verfallen um den "Wechselsatz" (1) der Statistik und dem Energiesatz
zu retten. Nämlich die Möglichkeit, es könnten elektrisch neutrale
Teilchen, die ich Neutronen nennen will, in den Kernen existieren,
welche den Spin 1/2 haben und das Ausschließungsprinzip befolgen und
sich von Lichtquanten ausserdem noch dadurch unterscheiden, dass sie
nicht mit Lichtgeschwindigkeit laufen. Die Masse der Neutronen
müsste von derselben Grossenordnung wie die Elektronenmasse sein und
jedemfalls nicht grösser als 0,01 Protonenmasse... Das kontinuierliche
beta-Spektrum wäre dann verständlich unter der Annahme, dass beim
Beta-Zerfall mit dem Elektron jeweils noch ein Neutron emittiert
wird, derart, dass die Summe der Energien von Neutron und Elektron
konstant ist.

Neary 1940

Experimental Validation

Detection of a neutrino from a nuclear reactor

Detection of the Free Neutrino*

F. REINES AND C. L. COWAN, JR.

Los Alamos Scientific Laboratory, University of California,
Los Alamos, New Mexico

(Received July 9, 1953; revised manuscript received September 14, 1953)

An experiment¹ has been performed to detect the free neutrino. It appears probable that this aim has been accomplished although further confirmatory work is in progress. The cross section for the reaction employed,

$$\nu - + p \rightarrow n + \beta^+, \quad (1)$$

has been calculated^{2,3} from beta-decay theory to be given by the expression,

$$\sigma = \left(\frac{G^2}{2\pi} \right) \left(\frac{\hbar}{mc} \right)^2 \left(\frac{p}{mc} \right)^2 \left(\frac{1}{v/c} \right), \quad (2)$$

where σ = cross section in barns; p , m , v = momentum, mass, and velocity of emitted positron (cgs units); c = velocity of light (cm/sec); $2\pi\hbar$ = Planck's constant (cgs units); and G^2 = dimensionless lumped β -coupling constant (= 55 from measurements of neutron and tritium β decay).⁴ An estimate of the fission frag-

Reines, Cowan 1953

A Brief History of Neutrino Anomalies

Experimental Anomaly


Theoretical Resolution

Experimental Validation


A Brief History of Neutrino Anomalies

Experimental Anomaly

Observed deficits of solar and atmospheric neutrinos



Theoretical Resolution




Experimental Validation

A Brief History of Neutrino Anomalies

Experimental Anomaly


Observed deficits of solar and atmospheric neutrinos



Theoretical Resolution

Oscillations* from mixing between mass and flavor states

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = \begin{pmatrix} U_{e1} & U_{e2} & U_{e3} \\ U_{\mu 1} & U_{\mu 2} & U_{\mu 3} \\ U_{\tau 1} & U_{\tau 2} & U_{\tau 3} \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}$$



$$P(\nu_\alpha \rightarrow \nu_\beta) \approx \sin^2 2\theta \sin^2 \left(\frac{\Delta m^2 L}{4E} \right)$$


*and matter effects

Experimental Validation

A Brief History of Neutrino Anomalies

Experimental Anomaly


Observed deficits of solar and atmospheric neutrinos



Theoretical Resolution

Oscillations* from mixing between mass and flavor states

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = \begin{pmatrix} U_{e1} & U_{e2} & U_{e3} \\ U_{\mu 1} & U_{\mu 2} & U_{\mu 3} \\ U_{\tau 1} & U_{\tau 2} & U_{\tau 3} \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}$$





$$P(\nu_\alpha \rightarrow \nu_\beta) \approx \sin^2 2\theta \sin^2 \left(\frac{\Delta m^2 L}{4E} \right)$$

*and matter effects

Experimental Validation

New measurements of solar, atmospheric, and reactor neutrinos



A Brief History of Neutrino Anomalies

Experimental Anomaly


Theoretical Resolution

Experimental Validation


A Brief History of Neutrino Anomalies

Experimental Anomaly

Excesses of ν_e and $\bar{\nu}_e$ events in short baseline neutrino experiments



Theoretical Resolution




Experimental Validation

A Brief History of Neutrino Anomalies

Experimental Anomaly

Excesses of ν_e and $\bar{\nu}_e$ events in short baseline neutrino experiments




Theoretical Resolution

Unresolved!

Unknown systematic uncertainties?

eV-scale sterile neutrinos?




Experimental Validation

GeV-scale heavy neutrinos?

A Brief History of Neutrino Anomalies

Experimental Anomaly


Excesses of ν_e and $\bar{\nu}_e$ events in short baseline neutrino experiments



Theoretical Resolution

Unresolved!

Unknown systematic uncertainties?



Experimental Validation




eV-scale sterile neutrinos?

GeV-scale heavy neutrinos?

A Brief History of Neutrino Anomalies

Experimental Anomaly


Excesses of ν_e and $\bar{\nu}_e$ events in short baseline neutrino experiments



Theoretical Resolution

Unresolved!

Unknown systematic uncertainties?



Experimental Validation




eV-scale sterile neutrinos?

GeV-scale heavy neutrinos?

This talk!



Outline

1. **MiniBooNE:** A long-standing neutrino anomaly
2. **Heavy neutrinos in plastic:** constraints on a promising MiniBooNE solution
3. **Heavy neutrinos in ice and water:** searches at neutrino telescopes
4. **Heavy neutrinos (and more) in water and rock:** new detectors for collider neutrinos



Outline

1. **MiniBooNE:** A long-standing neutrino anomaly
2. **Heavy neutrinos in plastic:** constraints on a promising MiniBooNE solution
3. **Heavy neutrinos in ice and water:** searches at neutrino telescopes
4. **Heavy neutrinos (and more) in water and rock:** new detectors for collider neutrinos





The MiniBooNE Experiment

- 800-ton CH_2 Cherenkov detector
- Situated along Fermilab's Booster Neutrino Beam
 - ~540 m from the beryllium target
- Observes mostly ν_μ and $\bar{\nu}_\mu$ from charged pion decays





FNAL booster
(8 GeV protons)




The MiniBooNE Anomaly

Electrons:
“fuzzy” rings from multiple scattering



4.8 σ excess of electron-like events




What could it be?

Cherenkov limitations:

- Electrons/photons indistinguishable
- No hadronic information

The excess could be...




What could it be?

Cherenkov limitations:

- Electrons/photons indistinguishable
- No hadronic information

The excess could be...

1. True electron neutrinos?



What could it be?


Cherenkov limitations:

- Electrons/photons indistinguishable
- No hadronic information

The excess could be...

1. True electron neutrinos?

$$\text{Recall } P(\nu_\mu \rightarrow \nu_e) \propto \sin^2(\Delta m^2 L / 4E)$$




What could it be?

Cherenkov limitations:

- Electrons/photons indistinguishable
- No hadronic information

The excess could be...

1. True electron neutrinos?



Recall $P(\nu_\mu \rightarrow \nu_e) \propto \sin^2(\Delta m^2 L / 4E)$

$$\Delta m_{23}^2 \sim 2 \times 10^{-3} \text{ eV}^2, E_\nu \sim 500 \text{ MeV} \implies L_{\text{osc}} \sim 200 \text{ km}$$

What could it be?

Cherenkov limitations:

- Electrons/photons indistinguishable
- No hadronic information


The excess could be...

1. True electron neutrinos?

Recall $P(\nu_\mu \rightarrow \nu_e) \propto \sin^2(\Delta m^2 L / 4E)$

$$\Delta m_{23}^2 \sim 2 \times 10^{-3} \text{ eV}^2, E_\nu \sim 500 \text{ MeV} \implies L_{\text{osc}} \sim 200 \text{ km}$$

$L_{\text{MB}} \sim 500 \text{ m}$, much too short for $\nu_\mu \rightarrow \nu_e$ oscillations via Δm_{23}^2 !



What could it be?


Cherenkov limitations:

- Electrons/photons indistinguishable
- No hadronic information

The excess could be...

1. True electron neutrinos?

What about extra intrinsic
electron neutrinos in the BNB?



What could it be?

Cherenkov limitations:



- Electrons/photons indistinguishable
- No hadronic information

The excess could be...

1. True electron neutrinos?

What about extra intrinsic
electron neutrinos in the BNB?

**MicroBooNE data disfavor this
hypothesis at the 3σ confidence level**



What could it be?

Cherenkov limitations:



- Electrons/photons indistinguishable
- No hadronic information

The excess could be...

1. True electron neutrinos? 

What about extra intrinsic
electron neutrinos in the BNB?

**MicroBooNE data disfavor this
hypothesis at the 3σ confidence level**




What could it be?

Cherenkov limitations:

- Electrons/photons indistinguishable
- No hadronic information

The excess could be...

1. True electron neutrinos? ~~X~~
2. Mis-modeled photon background?



What could it be?

Cherenkov limitations:


- Electrons/photons indistinguishable
- No hadronic information

The excess could be...



1. True electron neutrinos? ~~X~~

2. Mis-modeled photon background?

π^0 misidentification
background constrained
in-situ and disfavored by
the radial distribution



Example of a π^0 mis-ID



What could it be?


Cherenkov limitations:

- Electrons/photons indistinguishable
- No hadronic information

The excess could be...

1. True electron neutrinos? ~~X~~
2. Mis-modeled photon background?

Rare Δ decays to single photons are not constrained in situ by MiniBooNE



What could it be?

Cherenkov limitations:



- Electrons/photons indistinguishable
- No hadronic information

The excess could be...

1. True electron neutrinos? ~~X~~
2. Mis-modeled photon background?

Rare Δ decays to single photons are not constrained in situ by MiniBooNE

MicroBooNE data also disfavor a MiniBooNE-like excess of $\Delta \rightarrow N\gamma$ events at the 95% confidence level



What could it be?

Cherenkov limitations:



- Electrons/photons indistinguishable
- No hadronic information

The excess could be...

1. True electron neutrinos? X
2. Mis-modeled photon background? X

Rare Δ decays to single photons are not constrained in situ by MiniBooNE

MicroBooNE data also disfavor a MiniBooNE-like excess of $\Delta \rightarrow N\gamma$ events at the 95% confidence level



What could it be?

Cherenkov limitations:



- Electrons/photons indistinguishable
- No hadronic information

The excess could be...

1. True electron neutrinos? X
2. Mis-modeled photon background? X (See also Brdar, Kopp 2021)

Rare Δ decays to single photons are not constrained in situ by MiniBooNE

MicroBooNE data also disfavor a MiniBooNE-like excess of $\Delta \rightarrow N\gamma$ events at the 95% confidence level




What could it be?

Cherenkov limitations:

- Electrons/photons indistinguishable
- No hadronic information

The excess could be...

1. True electron neutrinos? X
2. Mis-modeled photon background? X
3. New physics?



What could it be?




Cherenkov limitations:

- Electrons/photons indistinguishable
- No hadronic information

The excess could be...

1. True electron neutrinos? X
2. Mis-modeled photon background? X
3. New physics?

An eV-scale sterile neutrino can induce
 $\nu_\mu \rightarrow \nu_e$ oscillations at short baselines



What could it be?


Cherenkov limitations:

- Electrons/photons indistinguishable
- No hadronic information

The excess could be...

1. True electron neutrinos? X
2. Mis-modeled photon background? X
3. New physics?

So why haven't we declared victory?



What could it be?

Cherenkov limitations:

- Electrons/photons indistinguishable
- No hadronic information

The excess could be...



1. True electron neutrinos?

2. Mis-modeled photon background?

3. New physics?

So why haven't we declared victory?

**Significant internal tension
in sterile neutrino global fits**



No positive evidence for ν_μ disappearance, up to a persistent $\sim 2\sigma$ hint from IceCube
PRL 133, 201804
(2024)

What could it be?

Cherenkov limitations:


- Electrons/photons
- No hadronic info

The excess could be


1. True electron neutrino
2. Mis-modeled photons
3. New physics?

So why haven't we declared victory?

**Significant internal tension
in sterile neutrino global fits**




None of our leading Standard Model or BSM hypotheses for the excess seem to work...
We need a new model




No positive evidence for ν_μ disappearance, up to a persistent $\sim 2\sigma$ hint from IceCube
PRL 133, 201804 (2024)

A hint for a two component solution




A hint for a two component solution

**A broad angle
component at
lower energies**



A hint for a two component solution


**A broad angle
component at
lower energies**



**A forward
component
that extends
to higher
energies**

Outline

1. **MiniBooNE:** A long-standing neutrino anomaly
2. **Heavy neutrinos in plastic:** constraints on a promising MiniBooNE solution
3. **Heavy neutrinos in ice and water:** searches at neutrino telescopes
4. **Heavy neutrinos (and more) in water and rock:** new detectors for collider neutrinos




Can we find a two component solution for MiniBooNE?

[1] [Hardin+ 2211.02610](#) [2] [Vergani, NK+ PRD 104, 095005](#) [3] [NK+ PRD 107, 055009](#)

Can we find a two component solution for MiniBooNE?


(1) An eV-scale sterile neutrino



[1] Hardin+ 2211.02610 [2] Vergani, NK+ PRD 104, 095005 [3] NK+ PRD 107, 055009


Can we find a two component solution for MiniBooNE?

(1) An eV-scale sterile neutrino



Subdominant contribution to the MiniBooNE excess while still explaining other short baseline neutrino anomalies [1,2]

(2) A dipole-portal heavy neutral lepton (HNL)




These new interactions explain the bulk of the MiniBooNE excess, relieving tension in sterile neutrino global fits ($4.8\sigma \rightarrow 2.3\sigma$) [2,3]

[1] Hardin+ 2211.02610 [2] Vergani, NK+ PRD 104, 095005 [3] NK+ PRD 107, 055009

Can we find a two component solution for MiniBooNE?

(1) An eV-scale sterile neutrino



Explaining the MiniBooNE excess through a mixed model of neutrino oscillation and decay


S. Vergani, N. W. Kamp, A. Diaz, C. A. Argüelles, J. M. Conrad, M. H. Shaevitz, and M. A. Uchida
Phys. Rev. D **104**, 095005 – Published 8 November 2021

Dipole-coupled heavy-neutral-lepton explanations of the MiniBooNE excess including constraints from MINERvA data

N. W. Kamp, M. Hostert, A. Schneider, S. Vergani, C. A. Argüelles, J. M. Conrad, M. H. Shaevitz, and M. A. Uchida
Phys. Rev. D **107**, 055009 – Published 9 March 2023

[1] Hardin+ 2211.02610 [2] Vergani, NK+ PRD 104, 095005



(2) A dipole-portal heavy neutral lepton (HNL)





Mixed Model Fit Strategy

1. Perform a global sterile neutrino fit without MiniBooNE to fix the sterile neutrino parameters [1]
2. Simulate HNL upscattering and decay in MiniBooNE using the open source SIREN simulation package [2]
3. Fit the remaining excess for the preferred HNL mass and dipole coupling

[1] [Vergani, NK+ PRD 104, 095005](#) [2] [A. Schneider, NK, A. Wen. 2024](#)







Dipole-portal HNLs @ MiniBooNE



HNLS in Plastic @ MINERvA

- Single showers from dipole-portal HNL decays would show up in MINERvA neutrino-electron elastic scattering measurements [1]
- We simulate upscattering and decay in a realistic MINERvA detector using SIREN





[1] MINERvA 2019

Kamp+ PRD 107, 055009

HNLs in Plastic @ MINERvA



- We use the high dE/dx sideband region to set constraints on dipole-portal HNLs
- Most stringent MINERvA constraints do not rule out the MiniBooNE-preferred region at the 95% CL





Kamp+ PRD 107, 055009


HNLs in Plastic @ MINERvA



- We use the high
- Most stringent N
- region at the 95%



one-portal HNLs
NE-preferred




Kamp+ PRD 107, 055009

HNLS in Plastic @ ND280(+)



- ND280's gaseous time projection chambers (TPCs) have low single shower backgrounds
- T2K leveraged this to search for e^+e^- pairs from mass-mixed HNL decays [1]
- **We repurpose their results to set constraints on dipole-portal HNLs**



[arXiv:2412.15051](https://arxiv.org/abs/2412.15051)

Constraints and Sensitivities for Dipole-Portal Heavy Neutral Leptons from ND280 and ND280+

M-S. Liu,^{1,*} N.W. Kamp,^{2,†} and C.A. Argüelles^{2,‡}

¹*Cavendish Laboratory, University of Cambridge, Cambridge, CB3 0HE, UK*

²*Department of Physics and Laboratory for Particle Physics and Cosmology, Harvard University, Cambridge, MA 02138, USA*



M-S. Liu

[1] [T2K 2019](#)


[M. Lamoureux thesis](#)

HNLs in Plastic @ ND280(+)




**3D rendering of our SIREN-based implementation
of the nominal and upgraded ND280 detector**

ND280










ND280+



M-S Liu, NK, C. Argüelles 2024


HNLs in Plastic @ ND280(+)



$$\Gamma_{\mathcal{N} \rightarrow \nu\gamma} \propto m_{\mathcal{N}}^3$$

M-S Liu, NK, C. Argüelles 2024

HNLS in Plastic @ ND280(+)



The 2019 T2K search observes zero e^+e^- pairs in the ND280 gas TPCs, constraining the region of parameter space preferred by MiniBooNE

The addition of three years of ND280 upgrade data will further improve the sensitivity

Caveat: these constraints assume the same efficiency for tagging mass-mixed and dipole-portal HNL decays

H

Takeaway:

+)




Dipole-portal HNLs are a promising explanation of the MiniBooNE excess, though they face constraints from MINERvA and ND280 data

observes zero
280 gas TPCs,
in of parameter

space preferred by MiniBooNE

The addition of three years of ND280 upgrade data will further improve the sensitivity

Caveat: these constraints assume the same efficiency for tagging mass-mixed and dipole-portal HNL decays



H


Takeaway:

+



Dipole-portal HNLs are a promising explanation of the MiniBooNE excess, though they face constraints from MINERvA and ND280 data

observes zero
280 gas TPCs,
in of parameter
space preferred by
MiniBooNE



However:



Single showers in MINERvA and ND280 are not “smoking-gun” evidence for dipole-portal HNLs. Can we make a more compelling search elsewhere?

ree years of
a will further
nsitivity



nts assume the
agging mass-
tal HNL decays

Outline

1. **MiniBooNE:** A long-standing neutrino anomaly
2. **Heavy neutrinos in plastic:** constraints on a promising MiniBooNE solution
3. **Heavy neutrinos in ice and water:** searches at neutrino telescopes
4. **Heavy neutrinos (and more) in water and rock:** new detectors for collider neutrinos





The IceCube Detector




Quanta 2023

IceCube Event Categories




"Through-going track"

ν_μ charged-current DIS outside the active volume



"Starting track"


ν_μ charged-current DIS inside the active volume





"Cascade"

$\nu_{e,\tau}$ charged-current DIS or ν_α neutral-current DIS inside the active volume

IceCube Event Categories



Through-going track





Earliest photons





Latest photons


Starting track




Cascade



HNLS in Ice @ IceCube




- We look for two isolated cascades from the HNL upscattering and decay
- **This is a smoking gun signature* of dipole-portal HNLs!**
- First proposed in Coloma+ (2017)




*Double cascades are also a signature of ν_τ interactions at much higher energies [IceCube 2020]




HNLS in Ice @ IceCube





- Upscattering and decay of HNLs simulated using SIREN
- Photon propagation and detector response simulated using internal IceCube software



Atmospheric Neutrino Flux





- Atmospheric neutrinos are our main $\nu_\mu/\bar{\nu}_\mu$ source to produce dipole-portal HNLs
- Flux prediction from DAEMONFLUX [1]



[1] Yañez+ 2023



HNLS in Ice @ IceCube




Number of double cascade from dipole-portal HNLs per year:

1. Passing IceCube's data filtration system
2. With cascade separation > 5 m

HNLS in Ice @ IceCube



23 DUs Deployed



31x 3" PMTs



43 cm

18 Jun 2024


KM3NeT/ORCA



115 Detection Units
18 DOMs / DU

200 m

7 Mton detector



6




We can also
search for HNLs
in KM3NeT

The longer photon
scattering length of
water v.s. ice is
advantageous for event
reconstruction


Reproduced from
Joao Coelho, Neutrino 2024

HNLS in Water @ KM3NeT




- We begin with a SIREN-based simulation of dipole-portal HNL double cascades in ORCA
- Graph neural networks appear to have strong separation capability between double cascade signals and single cascade backgrounds

Sensitivity studies underway



HNLS in Water @ KM3NeT





- We begin with a SIREN-based simulation of dipole-portal HNL double cascades in ORCA

-

Takeaway:



IceCube and KM3NeT are performing searches for the “smoking-gun” double cascade signature of dipole-portal HNLs



J. Prado





Outline

1. **MiniBooNE:** A long-standing neutrino anomaly
2. **Heavy neutrinos in plastic:** constraints on a promising MiniBooNE solution
3. **Heavy neutrinos in ice and water:** searches at neutrino telescopes
4. **Heavy neutrinos (and more) in water and rock:** new detectors for collider neutrinos



The Dawn of Collider Neutrino Physics


Unique sensitivity to TeV neutrinos and long-lived particles produced in the forward direction at the LHC



FASER Collab. 2023


FASER Collab. 2024

LHC Neutrinos pass through Lake Geneva




This enables the construction of large-scale lake-and-surface-based detectors that evade muon backgrounds from the p-p collision

Thanks to Benjamin Weyer and Albert De Roeck for discussions on the beam geometry



LHC Neutrinos pass through Lake Geneva



This enables the construction of large-scale lake-and-surface-based detectors that evade muon backgrounds from the p-p collision

Thanks to Benjamin Weyer and

arXiv:2501.03278

Lake- and Surface-Based Detectors for Forward Neutrino Physics

Nicholas W. Kamp,^{1,*} Carlos A. Argüelles,^{1,†} Albrecht Karle,^{2,‡} Jennifer Thomas,^{2,3,§} and Tianlu Yuan^{2,¶}

¹Department of Physics and Laboratory for Particle Physics and Cosmology, Harvard University, Cambridge, MA 02138, US

²Department of Physics and Wisconsin IceCube Particle Astrophysics Center,

University of Wisconsin–Madison, Madison, WI 53706, USA

³Department of Physics and Astronomy, University College London, London, WC1E 6BT, UK

(Dated: January 14, 2025)



J. Thomas



A. Karle



C. Argüelles




T. Yuan


Distance from CMS Interaction Point [m]

Distance from LHCb Interaction Point [m]




SINE: Surface-based Integrated Neutrino Experiment



Signal definition: up-going muons from neutrino interactions in bedrock





UNDINE: UNDerwater Integrated Neutrino Experiment



NK+ 2025

LHC Forward Neutrino Flux

We use github.com/makelat/forward-nu-flux-fit for simulated forward neutrino flux samples




Event Rates

- We simulate DIS neutrino interactions along the LHCb and CMS beam using SIREN
- Cherenkov detectors enable flavor identification in UNDINE

These detectors offer a cost-effective opportunity to collect large samples of collider neutrino interactions


Dataset	Total	π, K	D, Λ_c
SINE (CC $\nu_\mu + \bar{\nu}_\mu$)	$10^{6.98}$	$10^{6.84}$	$10^{6.40}$
UNDINE (CC $\nu_e + \bar{\nu}_e$)	$10^{4.68}$	$10^{4.32}$	$10^{4.42}$
UNDINE (CC $\nu_\mu + \bar{\nu}_\mu$)	$10^{5.27}$	$10^{5.20}$	$10^{4.41}$
UNDINE (CC $\nu_\tau + \bar{\nu}_\tau$)	$10^{3.07}$	0	$10^{3.07}$
UNDINE (NC $\nu_\alpha + \bar{\nu}_\alpha$)	$10^{4.87}$	$10^{4.76}$	$10^{4.24}$



What can we do with
~a million collider
neutrinos?

Neutrino Cross Sections

- FASER recently reported first measurements of the total neutrino cross section at TeV energies [1,2]
- SINE and UNDINE can make complimentary measurements
- **Few-percent-level uncertainties with full dataset**
 - Comparable to theoretical uncertainty [3]



[1] FASER Collab. 2024


[2] FASER Collab. 2024

[3] Weigel+ 2024

NK+ 2025

Forward Charm Production in p-p Collisions

- Increasing forward charm production rates corresponds to...
 - 1. More high-energy muon neutrinos**
 - 2. More electron and tau neutrinos**
- Ratio measurements can distinguish between charm production models after 1 year
- Important implications for **intrinsic charm content of the proton [1]** and the **prompt atmospheric neutrino flux [2]**





[1] Maciula+ 2022

[2] Jeong+ 2023


NK+ 2025

Cosmic Muon Puzzle

- Excess of muons observed in cosmic ray air showers [1]
- **Hypothesis:** swapping probability f_s between pions and kaons in hadronic showers [2]
- SINE and UNDINE have complementary sensitivity to future FPF experiments [3]




HNLS in Rock and Water



- **Mass-mixed HNLs:** minimal extension of the Standard Model with HNLs
- Famously appear in the See-saw mechanism for neutrino mass generation
- Each neutrino flavor state α couples to the HNL by a small mixing $U_{\alpha 4}$


Can we look for them in SINE and UNDINE?





$$\mathcal{L} = \mathcal{L}_{SM} + i\bar{N}_I \gamma^\mu \partial_\mu N_I - F_{\alpha I} \bar{L}_\alpha \tilde{\Phi} N_I - \frac{M_{IJ}}{2} \bar{N}_I^c N_J + h.c.,$$

Abudllahi+ 2022


HNLS in Rock and Water



- Two ideas to look for mass-mixed HNLs in SINE and UNDINE




HNLS in Rock and Water




- Two ideas to look for mass-mixed HNLs in SINE and UNDINE

1. $m_{\mathcal{N}} \lesssim 10 \text{ GeV}$: Delayed muons with respect to the beam trigger from HNL time-of-flight





HNLS in Rock and Water




- Two ideas to look for mass-mixed HNLs in SINE and UNDINE

1. $m_{\mathcal{N}} \lesssim 10 \text{ GeV}$: Delayed muons with respect to the beam trigger from HNL time-of-flight
2. $m_{\mathcal{N}} > 10 \text{ GeV}$: Di-muons from $\mathcal{N} \rightarrow \mu(W^{(\star)} \rightarrow \mu\nu_\mu)$ or $\mathcal{N} \rightarrow \nu(Z^{(\star)} \rightarrow \mu\mu)$





HNLS in Rock and Water




- Two ideas to look for mass-mixed HNLs in SINE and UNDINE

1. $m_{\mathcal{N}} \lesssim 10 \text{ GeV}$: Delayed muons with respect to the beam trigger from HNL time-of-flight
2. $m_{\mathcal{N}} > 10 \text{ GeV}$: Di-muons from $\mathcal{N} \rightarrow \mu(W^{(\star)} \rightarrow \mu\nu_\mu)$ or $\mathcal{N} \rightarrow \nu(Z^{(\star)} \rightarrow \mu\mu)$

Sensitivity studies in progress



SINE Cosmic Backgrounds



Signal



- True coincidence from a ν -induced muon
- ~11 mHz per surface detector during HL-LHC

Background

- True coincidence from a single cosmic muon
- Accidental coincidence from two cosmic muons

*Water overburden reduces cosmic backgrounds in lake detector

SINE Cosmic Backgrounds




We use EcoMug to generate cosmic muons in a cylinder surrounding one of the shipping containers


Cosmic muon true coincidence rate: 1.67 kHz
Cosmic muon single panel rate: 1.62 kHz

Four Strategies for Background Rejection


1. Timing with respect to proton collision





2. Time difference between scintillator panels








3a. Up-going spatial information



3b. Two-dimensional spatial information




SINE Background Summary




Prototype SINE Detector

lhccommissioning.web.cern.ch/schedule/LHC-long-term.htm




**What can we do here with one
shipping crate detector?**

Shipping Container




Prototype SINE Detector



Ihc-commissioning



Shutdown/Technical stop
Protons physics
Ions (tbc after LS4)
Commissioning with beam
Hardware commissioning



Takeaway:
SINE and UNDINE are cost-effective experiments that can:
(1) collect large samples of collider neutrino interactions, and
(2) potentially set strong constraints on mass-mixed HNLs



Conclusion

Conclusion

The 4.8σ excess of electron-like events in MiniBooNE remains unexplained

Conclusion

The 4.8σ excess of electron-like events in MiniBooNE remains unexplained

Dipole-portal HNLs offer a promising explanation for the bulk of the excess

Conclusion

The 4.8σ excess of electron-like events in MiniBooNE remains unexplained

Dipole-portal HNLs offer a promising explanation for the bulk of the excess

MINERvA and ND280 data constrain dipole-portal HNLs

Conclusion

The 4.8σ excess of electron-like events in MiniBooNE remains unexplained

Dipole-portal HNLs offer a promising explanation for the bulk of the excess

MINERvA and ND280 data constrain dipole-portal HNLs

IceCube and KM3NeT are performing complimentary searches for the “smoking-gun” double cascade signature of dipole-portal HNLs

Conclusion

The 4.8σ excess of electron-like events in MiniBooNE remains unexplained

Dipole-portal HNLs offer a promising explanation for the bulk of the excess

MINERvA and ND280 data constrain dipole-portal HNLs

IceCube and KM3NeT are performing complimentary searches for the “smoking-gun” double cascade signature of dipole-portal HNLs

SINE and UNDINE can collect large samples of collider neutrino interactions, constraining Standard Model neutrino physics and potentially HNLs

Thank you!

Backups